

X-ray emission from the young brown dwarfs of the Taurus Molecular Cloud

N. Grosso¹, K. R. Briggs², M. Güdel², S. Guieu¹, E. Franciosini³, F. Palla⁴,
C. Dougados¹, J.-L. Monin^{1,5}, F. Ménard¹, J. Bouvier¹, M. Audard^{6,7}, and A. Telleschi²

¹ LAOG, UJF, CNRS, 414 rue de la Piscine, 38041 Grenoble, France

² Paul Scherrer Institut, 5232 Würenlingen und Villigen, Switzerland

³ INAF - OAPa, Piazza del Parlamento 1, 90134 Palermo, Italy

⁴ INAF - Osservatorio Astrofisico di Arcetri, Largo Enrico Fermi 5, 50125 Firenze, Italy

⁵ Institut Universitaire de France

⁶ Columbia Astrophysics Laboratory, 550 West 120th Street, New York, NY 10027, USA

⁷ Observatory of Geneva, 51 Chemin des Maillettes, 1290 Sauverny, Switzerland

Abstract. We surveyed with 15 *XMM-Newton* pointings, supplemented with one archival *Chandra* observation, 17 young brown dwarfs (BDs) with late M spectral types in the Taurus Molecular Cloud. Half of this sample (9 out of 17 BDs) is detected; 7 BDs are detected here for the first time in X-rays. We confirm several previous findings on BD X-ray activity: a log-log relation between X-ray and bolometric luminosity for stars and BDs detected in X-rays; a shallow log-log relation between X-ray fractional luminosity and mass; a log-log relation between X-ray fractional luminosity and effective temperature; a log-log relation between X-ray surface flux and effective temperature. We find no significant log-log correlation between the X-ray fractional luminosity and EW(H α). Accreting and nonaccreting BDs have a similar X-ray fractional luminosity. The X-ray fractional luminosity declines from low-mass stars to M-type BDs and, as a sample, the BDs are less efficient X-ray emitters than low-mass stars. We thus conclude that while the BD atmospheres observed here are mostly warm enough to sustain coronal activity, a trend is seen that may indicate its gradual decline due to the drop in photospheric ionization degree.

Key words. Stars: low-mass, brown dwarfs – X-rays: stars – ISM: individual objects: the Taurus Molecular Cloud

1. Introduction

The *XMM-Newton Extended Survey of the Taurus Molecular Cloud* (XEST; Güdel et al. 2007) is a large program designed to investigate the X-ray properties of young stellar/substellar objects in the Taurus Molecular Cloud (TMC). In particular, the area surveyed by 15 (out of 27) *XMM-Newton* pointings, supplemented with one archival *Chandra* observa-

tion, allows us to study 17 young (~ 3 Myr) brown dwarfs (BDs) with late M spectral types (Fig. 1; Grosso et al. 2007).

2. X-ray properties of the XEST BDs

We detected half of this sample (9 out of 17 BDs); 7 BDs are detected here for the first time in X-rays. We detected mainly the BDs with




Fig. 1. The *XMM-Newton* Extended Survey of the Taurus Molecular Cloud (XEST). Contours show the ^{12}CO emission of the TMC (Dame et al. 1987), overlaid on a visual extinction map (Dobashi et al. 2005). The 27 *XMM-Newton* fields of view of the XEST are plotted with continuous and dashed (archive) circles. The two squares near labels 3 and 8 show the archival *Chandra* fields of view also used in this work. Other bold lines outline areas surveyed for BDs with the *CFHT* (Guieu et al. 2006). This region hosts 42 young BDs of the TMC (Briceño et al. 1998, 2002; Luhman 2000, 2004, 2006; Martín et al. 2001; Guieu et al. 2006). Black dots indicate the 25 BDs of this region not surveyed with the XEST. The 17 BDs included in the present X-ray study are plotted with white dots (8 BDs not detected) and crosses (9 BDs detected), with labels referring to BD names.

luminosities greater than $\sim 0.01 L_{\odot}$, i.e. with spectral type M8 or earlier.

Considering only the BDs and low-mass (proto)stars detected in the XEST (Fig. 2), we determined a log-log regression fit between X-ray and bolometric luminosity, which is consistent with $\langle \log(L_X/L_*) \rangle = -3.5 \pm 0.4$. Including the 8 upper limits of undetected BDs, the median of $\log(L_X/L_*)$ is -4.0 for the XEST BDs.

The X-ray fractional luminosity of XEST BDs is hence lower than the one of XEST low-mass stars. To investigate the relation between X-ray fractional luminosity and physical parameters when one moves from low-mass stars to the substellar regime, we focus on objects with spectral type M0 or later. A shallow relation is found between the X-ray fractional luminosity and the mass. We find a relation between




Fig. 2. X-ray luminosity vs. bolometric luminosity for the BDs (diamonds), low-mass stars and protostars (white and black dots) of the XEST (Güdel et al. 2007). There are only a few low-mass stars not detected in the XEST. The dotted lines indicate logarithmic X-ray fractional luminosities of -5, -4, -3, -2.

the X-ray fractional luminosity and the effective temperature of our sample ranging from ~ 3840 K to ~ 2500 K, implying a decrease of the X-ray fractional luminosity by a factor of about 3 from hot photospheres of solar-mass stars to cooler atmospheres of M9 BDs.

We compare the X-ray fractional luminosities of BDs with field stars with spectral type M5V or later, which also show $\log(L_X/L_*)$ as high as ~ -3 to ~ -4 . Field stars with spectral type M7V and an age of 1 Gyr are twice as massive as a typical TMC BD with an M7 spectral type. Moreover, such very cool stars also have surface gravities about 40 times higher than in a typical young BD. This shows that the X-ray activity of BD coronae is not strongly dependent of the BD mass and surface gravity. The more fundamental relation that we found between X-ray activity and effective temperature agrees with the overall statistics for field dwarfs and BDs: of the 15 objects with spectral type M8.5 or later, only 4 have any detected quiescent X-ray emission; the rest are either not detected at all or (in 3 cases) detected only during strong flares.

We find no significant log-log correlation between the X-ray fractional luminosity and $\text{EW}(\text{H}\alpha)$. Accreting and nonaccreting BDs in the TMC have a similar X-ray fractional luminosity.

3. Conclusion

We confirm that there is no dramatic change of the magnetic activity at the stellar/substellar boundary. Young BDs of spectral type M are sufficiently warm to sustain an active corona. The X-ray surface flux (i.e., the X-ray luminosity divided by the stellar surface) decreases with the effective temperature, but the young BDs in the TMC have on average an X-ray surface flux which is still 7 times higher than the one observed in the solar corona at the solar cycle maximum.

Deeper X-ray observations of the coolest M-type BDs in the TMC are needed to investigate a possible turn-over of the fractional X-ray luminosity of TMC BDs around spectral type M9.

Acknowledgements. We thank the International Space Science Institute in Bern for significant financial support of the project team. This research is based on observations obtained with *XMM-Newton*, an ESA science mission with instruments and contributions directly funded by ESA Member States and NASA. X-ray astronomy research at PSI has been supported by the Swiss National Science Foundation (grants 20-66875.01 and 20-109255/1). M.A. acknowledges support from NASA grant NNG05GF92G, and a Swiss National Science Foundation Professorship (PP002-110504).

References

- Briceño, C. et al. 1998, *AJ*, 115, 2074
- Briceño, C. et al. 2002, *ApJ*, 580, 317
- Dame, T. M. et al. 1987, *ApJ*, 322, 706
- Dobashi, K. et al. 2005, *PASJ*, 57, 1
- Grosso, N. et al. 2007, *A&A*, in press
- Güdel, M. et al. 2007, *A&A*, in press
- Guieu, S. et al. 2006, *A&A*, 446, 485
- Luhman, K. L. 2000, *ApJ*, 544, 1044
- Luhman, K. L. 2004, *ApJ*, 617, 1216
- Luhman, K. L. 2006, *ApJ*, 645, 676
- Martín, E. L. et al. 2001, *ApJ*, 561, L195