

Understanding intrinsic galaxy alignments

1 Abstract

The apparent distortions and alignments of faint galaxy images due to the gravitational lensing effect by the large-scale structure of the Universe is one of the potentially most powerful probes of cosmology. This so-called cosmic shear effect maps the matter distribution at different epochs and is thus sensitive to both the expansion history and the growth of structure. Forthcoming cosmic shear surveys such as the ESA Euclid mission will enable unprecedented insight into the properties of dark matter and dark energy, and test the validity of general relativity on cosmological scales.

The anticipated high-quality measurements of cosmic shear require an accurate understanding and treatment of systematic effects. The major astrophysical effect that contaminates the cosmic shear signal is the intrinsic alignment of galaxies. During their formation and evolution, galaxies are subjected to tidal forces exerted by the surrounding large-scale structure, which causes the galaxies to align with the matter distribution and neighbouring galaxies. These alignments can exactly mimic the gravitational lensing signal and have been demonstrated to potentially yield a systematic signal that is up to two orders of magnitude above the statistical errors expected for Euclid. To date, observational constraints are weak and models simplistic, so that a major effort is needed to ensure the success of the Euclid mission and other cosmic shear surveys.

The planned workshops will bring together leading observational, theoretical, and numerical expertise on intrinsic alignments, with the goal to decisively advance the understanding of this astrophysical effect. The international team, whose core will be formed by members of the Euclid Intrinsic Alignments Tiger Team, is going to write a review on the state of the art of intrinsic alignment research and will construct and test a new comprehensive, physically motivated intrinsic alignment model. The results will then be used to update the Euclid survey strategy, identify what external calibration data needs to be acquired, and optimise the treatment of intrinsic alignments in the Euclid analysis pipeline.

By virtue of the infrastructure and financial support of ISSI, the team is going to advance the knowledge of intrinsic alignments and disseminate its findings to the community via a review, peer-reviewed articles, and technical documentation in a timely and critical step towards the reliable interpretation of the forthcoming cosmic shear measurements.

2 Scientific rationale

The small, coherent distortions of faint galaxy images induced by the gravitational lensing of the intervening large-scale structure constitute potentially one of the most powerful probes of cosmology (Albrecht et al. 2006, Dark Energy Task Force Report, astro-ph/0609591), with a signal that in principle is straightforward to interpret physically (Bartelmann & Schneider 2001, Phys. Reports, 340, 291). This so-called cosmic shear effect allows for a statistical analysis of the properties of the full matter distribution at different epochs, as well as of the spatial geometry of the Universe. Therefore it is capable of answering some of the key open questions in fundamental physics: what are the properties of dark matter, what is the explanation for the accelerated expansion of the cosmos, and is general relativity the correct theory of gravity on cosmological scales.

Consequently, cosmic shear has become one of the main drivers of a new generation of large imaging surveys including the Dark Energy Survey (DES, <http://www.darkenergysurvey.org>), the Hyper Suprime-Cam Survey (HSC, <http://www.naoj.org/Projects/HSC/index.html>), and the Kilo Degree Survey (KiDS, <http://www.astro-wise.org/projects/KIDS>). Cosmic shear will constitute one of the two primary probes of the ESA Cosmic Vision mission Euclid (<http://www.euclid-ec.org>; Laureijs et al. 2011, Euclid Assessment Study Report for the ESA Cosmic Visions, astro-ph/0912.0914), to be launched in 2020. Over the course of six years Euclid will survey the full extragalactic sky in the optical and near-infrared with a 1.2m telescope, with a strong focus on superior image quality and excellent redshift information for about 2×10^9 galaxies.

The resulting sub-percent precision on the cosmic shear signal anticipated for Euclid needs to be matched with equally outstanding constraints on potential systematic effects. The main astrophysical systematics affecting cosmic shear are generated by the intrinsic alignment of galaxies. The physical alignment of galaxy shapes mimics the apparent correlations of the shapes of galaxy images created by




Figure 1: Illustration of intrinsic galaxy alignments, as seen along the line of sight (left panels) and on the sky (right panels). *Top row:* The intrinsic shapes of galaxies become correlated with each other when they are subjected to tidal forces exerted by the same matter structures. For this to happen, galaxies have to be close both in redshift and on the sky. Due to this so-called II effect galaxy shapes tend to point towards matter overdensities and become collinear, i.e. positively correlated. *Bottom row:* Alignments of galaxy shapes are also observed if a matter structure aligns a nearby galaxy and at the same time contributes to the lensing distortion of the image of a galaxy in the background. While the foreground galaxy tends to point towards matter overdensities, the lensing distortion of the background object is preferentially tangential around mass concentrations, so that this correlation, called GI effect, is negative. (Bottom right panel from Hirata & Seljak 2004)

the gravitational lensing effect. Thus it is vital to understand the intrinsic alignment signal and separate it from cosmic shear in order to extract the cosmological information in an unbiased way. Provided the separation of the signals works, the high-precision measurements achievable with the forthcoming surveys will also put tight constraints on intrinsic alignment models, thereby converting intrinsic alignments from a mere nuisance signal to an independent and valuable astrophysical probe of the role of the large-scale environment in galaxy formation and evolution.

However, the current knowledge of the physics behind intrinsic galaxy alignments is poor, threatening the successful cosmological exploitation of the forthcoming large imaging surveys. Surveys like DES, HSC, and KiDS are already taking data, or will shortly go into survey mode, and will yield a statistical accuracy on cosmic shear measurements that requires considerable improvement in the calibration of intrinsic alignments on timescales of not much more than a year from now. In the same period any calibration samples or high-level requirements that are needed to guarantee the robust calibration of Euclid cosmic shear measurements must be identified in order to be integrated into the implementation phase in time.

2.1 The current knowledge of alignment processes

If galaxies are subjected to tidal fields generated by their large-scale environment during formation and evolution, it is expected that their shapes (e.g. measured via the quadrupole of the light distribution) become correlated with the large-scale gravitational field (see Fig. 1 for an illustration). Then the intrinsic shapes of two physically close galaxies become aligned if they are influenced by the same gravitational forces (called the II effect). If a certain matter distribution aligns a nearby galaxy and at the same time contributes to the lensing distortion of a background galaxy image, one obtains correlations between gravitational distortion and intrinsic shape (called the GI effect).

In a simple analytical ansatz, referred to as the linear alignment model, it is assumed that the (projected) ellipticity of early-type galaxies is proportional to the quadrupole of the gravitational field at the time of the galaxy's formation (e.g. Hirata & Seljak 2004, PRD, 70, 063526). For late-type galaxies the alignment is assumed to be governed by their angular momentum, which in turn is quadratic in the gravitational potential according to tidal torquing theory (Peebles 1969, ApJ, 155, 393). This quadratic alignment model predicts a null signal to first order, while the linear alignment model results in two-point statistics for the II and GI signals that are rescaled versions of the linear matter power spectrum.

The linear alignment model has been phenomenologically extended by using the non-linear matter power spectrum instead of the linear one to account for small-scale effects (Bridle & King 2007, NJPh, 9,

444), or adding luminosity or extra redshift dependencies (see e.g. Joachimi et al. 2011, A&A, 527, 26). While these models are found to be in good agreement with current observations (Joachimi et al. 2011, A&A, 527, 26; Heymans et al. 2013, astro-ph/1303.1808), they lack physical motivation beyond the basic linear alignment model.

Schneider & Bridle (2010, MNRAS, 402, 2127) made a first attempt at a more consistent description of intrinsic alignments on small scales by devising a halo model. **The current implementation of the intrinsic alignment halo model simplistically assumes that the shapes of satellite galaxies all point towards the centre of the host halo and misses essential dependencies such as those on galaxy type and colour. However, the halo model can readily be extended to incorporate these and other more complex features, providing a clear framework to identify the free ingredients of the model (such as luminosity functions or radial alignment profiles of satellite galaxies) and how they can be constrained observationally.**

Complementary to the halo model ansatz, one can study intrinsic alignment signals in simulations. However, the accuracy which can eventually be achieved in this approach is currently unknown. Hydrodynamic simulations with a dark matter and baryon component could in principle produce realistic galaxy shapes and alignments, but their results depend strongly on the input sub-grid physics (see e.g. Semboloni et al. 2011, MNRAS, 417, 2020) and it is unclear if and when computational power will allow us to run sufficiently large simulations to beat down sample variance on intrinsic alignment signals. It will be possible to run hierarchical suites of dark matter-only simulations which jointly reach the required cosmological volume and mass resolution, but there remains a lot of freedom in modelling the link between dark matter halo shapes and alignments and the corresponding galaxy properties (Joachimi et al. 2013, astro-ph/1203.6833).

2.2 Observational constraints on intrinsic alignments

Due to the degeneracy with the cosmic shear signal, intrinsic alignments are difficult to constrain directly via observations. Measurements of the Π signal are restricted to low redshifts where the contribution by gravitational lensing is negligible (e.g. Brown et al. 2002, MNRAS, 333, 501). Constraints on the GI term are inferred from measurements of how strongly the orientation of intrinsic shapes of galaxies is aligned with the large-scale distribution of galaxies. This requires a translation from the clustering of galaxies to the clustering of the underlying dark matter distribution, at present done via the assumption of linear galaxy bias (see Mandelbaum et al. 2006, MNRAS, 367, 611). This step limits the interpretation of measurements to relatively large scales (comoving galaxy separations above $\sim 10 \text{ Mpc}/h$). A physically motivated and sufficiently complex halo model could simultaneously predict the clustering signal of relevant galaxy samples and hence remove this barrier.

To date detections have mostly been limited to luminous early-type samples (Mandelbaum et al. 2006, MNRAS, 367, 611; Hirata et al. 2007, MNRAS, 381, 1197; Joachimi et al. 2011, A&A, 527, 26), covering redshifts out to $z \sim 0.7$. Mandelbaum et al. (2011, MNRAS, 410, 844) analysed the largest late-type galaxy sample hitherto based on WiggleZ redshifts, but do not detect intrinsic alignments largely due to very large uncertainties induced by the low number density of the sample. On small scales the alignment of satellite galaxy shapes in galaxy clusters have been studied, but so far detections are tentative and partially in conflict with each other (see e.g. Hao et al. 2011, ApJ, 740, 39; Hung & Ebeling 2012, MNRAS, 421, 3229).

A typical galaxy in the Euclid cosmic shear survey for which intrinsic alignments will need to be well understood will be a quiescent late-type galaxy around redshifts of unity. **In the parameter space spanned by redshift, luminosity, and galaxy type we are currently far from constraining intrinsic alignments observationally among galaxies typically analysed with Euclid. Forthcoming surveys will explore much larger portions of this parameter space but will still find it hard to yield tight direct constraints on the intrinsic alignment signals relevant for Euclid. Therefore it is paramount to be able to rely on a robust physical intrinsic alignment model which can be extrapolated to the faint high-redshift objects that dominate the Euclid galaxy samples.**

Any future observations that will help improving our understanding of intrinsic alignments require top-quality imaging combined with precise redshift information (either spectroscopic or very precise photometric redshifts from multi-band photometry; see e.g. the proposed PAU Survey, <http://www.pausurvey.org>) for all galaxies. The resulting galaxy sample needs to cover significant cosmological volumes to beat down sample variance, and has to be reasonably dense on the sky to suppress shot noise in the er-

ror budget. Such tight requirements require careful and early planning with regards to making sure that suitable planned surveys take intrinsic alignments into consideration as a major science case.

2.3 Intrinsic alignment mitigation strategies in cosmological weak lensing

Both observations and simulations suggest that for certain galaxy samples intrinsic alignments could contribute up to 10% to the correlation signal of galaxy shapes (Heavens et al. 2000, MNRAS, 319, 649; Heymans et al. 2006, MNRAS, 371, 750; Mandelbaum et al. 2006, MNRAS, 367, 611), so that ignoring intrinsic alignments would have a devastating effect on imminent and future cosmic shear analyses. All forthcoming surveys will obtain good photometric redshifts for each galaxy which can be used to isolate intrinsic alignment signals to some extent without further knowledge about the underlying physical processes. The II signal is restricted to physically close pairs of galaxies (since the galaxies have to be subjected to the same tidal gravitational field) and is thus straightforward to suppress (King & Schneider 2002, A&A, 396, 411). The GI term has a contribution from gravitational lensing and therefore a scaling with redshift that is very similar to cosmic shear. Nonetheless it is possible to remove the GI signal, but one can only do so robustly with a substantial loss in cosmological constraining power which is unacceptable for surveys like Euclid (Joachimi & Schneider 2008, A&A, 488, 829; Joachimi & Schneider 2009, A&A, 507, 105).

Further proposed mitigation techniques exploit information from polarised emission from galaxies (currently tested in SuperCLASS, <http://www.e-merlin.ac.uk/legacy/projects/superclass.html>) or spatially resolved radial velocity maps. These quantities are unaffected by gravitational lensing and can be used to extract the intrinsic galaxy orientation (Brown & Battye 2011, MNRAS, 410, 2057; Morales 2006, ApJ, 650, 21). These methods would only be directly applicable to radio observations (e.g. with the SKA), but could help calibrate intrinsic alignments in optical data that overlaps with radio surveys.

A promising way to self-calibrate intrinsic alignments, in particular the GI term, with data accessible from a cosmic shear survey alone was investigated by Bernstein (2009, ApJ, 695, 652) and Joachimi & Bridle (2010, A&A, 523, 1). The method takes advantage of the galaxy clustering information that comes for free with cosmic shear data. This can be used to measure galaxy bias as well as the alignment of galaxy shapes towards galaxy overdensities, which yields an estimate of the GI contribution. Joachimi & Bridle evaluated the performance of the self-calibration technique (see also Kirk et al. 2012, MNRAS, 424, 1647), choosing a minimum-knowledge ansatz by parametrising intrinsic alignments and galaxy bias as a grid flexible in both redshift and spatial scales, with more than a hundred nuisance parameters in total. Even so one can recover the cosmological information provided by a pure cosmic shear signal to a good degree.

This approach has subsequently been applied in forecasts of the performance of the Euclid mission, impacting on the top-level cosmological constraints achievable with Euclid, as well as the requirements on photometric redshift measurements, which are driven by the need to separate intrinsic alignments and cosmic shear (Laureijs et al. 2011, astro-ph/0912.0914). For a given amount of observation time, a medium-deep survey with maximal survey area optimises cosmological constraints achievable with cosmic shear (Amara & Refregier 2007, MNRAS, 381, 1018). Including intrinsic alignments drives the optimum towards a deeper and smaller survey because of the need for a long baseline in redshift which again helps separating the intrinsic alignment and cosmic shear signals (Laureijs et al. 2011, astro-ph/0912.0914).

However, the general grid parametrisation may be too flexible and involves fairly arbitrary choices (such as the number of grid nodes) about the degree of flexibility. Moreover it is difficult to link the nuisance parameters of this parametrisation to physical observables, so that prior constraints are challenging to incorporate. **Replacing the minimum-knowledge approach for intrinsic alignments in Euclid performance forecasts by a robust, physically motivated intrinsic alignment model will minimise the number of ill-motivated choices in the modelling and dramatically simplify the identification and inclusion of prior information on the model. Consequently, key science requirements for Euclid as well as its survey and analysis strategy will be verified or updated with much improved confidence.**

Furthermore, the alignment mechanisms of a well-working halo model could readily be used to create mock cosmic shear surveys from large suites of N-body simulations that include a realistic intrinsic alignment signal. This will be paramount for the central task of developing and optimising intrinsic alignment mitigation and calibration techniques to be employed in the Euclid cosmic shear data analysis.

3 Scope of the project

The goal of this proposal is to bring together an international team of experts on intrinsic galaxy alignments who are going to

1. gain an overview and discuss in detail the current state of the art in the understanding of intrinsic alignments, including modelling attempts, results from simulations and observations, and mitigation strategies in cosmic shear analyses;
2. jointly work towards a comprehensive halo model of intrinsic alignments, identifying key parameters and ways to constrain them observationally;
3. review the Euclid science requirements impacted on by intrinsic alignments, repeat cosmological forecasts with the updated intrinsic alignment model, and update requirements, survey strategy, and analysis methods if needed.

The two proposed meetings at ISSI Bern constitute the central part of this effort where the work is initiated, organised, and reviewed. It is expected that the project will yield the following outcomes:

1. an extensive review paper on the current knowledge of intrinsic alignments;
2. a novel, comprehensive, and physically motivated intrinsic alignment halo model, tested against current observations (resulting in at least one refereed publication);
3. a road map to constrain free parameters of the intrinsic alignment model observationally, including the planning of survey involvement and/or preparation of proposals;
4. an update/verification of Euclid science requirements, survey strategies, and analysis methods (one refereed publication).

These results will mark critical and timely steps towards a successful cosmological exploitation of Euclid as well as the forthcoming ground-based imaging surveys because, as outlined above, significant progress in the understanding of intrinsic alignments is required on time scales of about a year.

4 Proposal details

4.1 Schedule

We plan to have a kick-off meeting before the end of the year, ideally in autumn, to review the state of the art in intrinsic alignment research and identify the tasks to be worked on in the following months. This will be followed by a second meeting after 6 to maximally 12 months, where the results of our work will be presented and steps towards finalising the projects, such as writing up papers, be undertaken. Both meetings are planned to last 5 days. All team members are committed to attending both meetings, should academic duties such as teaching obligations allow.

4.2 Facilities

We will require standard facilities that should readily be available at ISSI. This includes two meeting rooms (to allow for work being carried out in parallel) equipped with projectors, plenty of white board space, and a stable, fast WiFi connection. Power adaptors to the Swiss system will also be needed for team members' laptops.

4.3 Financial support

We apply for the standard financial support offered by ISSI, covering a per diem for the team members during their stay at ISSI and travel costs for the team leader. We will also benefit from the Young Scientist scheme by requesting funding for one or two senior PhD students or junior postdocs important to the success of our project.

4.4 Added value

The role of ISSI will be critical to the success of the proposed project via its financial and in-kind support that enables two focused week-long meetings of international experts in intrinsic galaxy alignments. We will have the opportunity to meet in an inspiring location and atmosphere, avoiding the usual distractions faced at institutes or during conferences.

5 Team members

1. **Benjamin Joachimi** (*coordinator*): IfA, University of Edinburgh, Royal Observatory, Blackford Hill, Edinburgh EH9 3HJ, United Kingdom; phone: +44-131-6688351; fax: +44-131-6688416; email: bj@roe.ac.uk
2. **Christopher Bonnet**: Institut de Ciències de l'Espai, CSIC/IEEC, Fac de Ciències, Torre C5 par-2, Barcelona 08193, Spain; phone: +34-935-868042; fax: N/A; email: c.bonnett@gmail.com
3. **Michael Brown**: Jodrell Bank Centre for Astrophysics, School of Physics & Astronomy, University of Manchester, Oxford Road, Manchester M13 9PL, UK; phone: +44-161-3063913; fax: +44-161-2754247; email: mbrown@jb.man.ac.uk
4. **Marcello Cacciato**: Leiden Observatory, Leiden University, Niels Bohrweg 2, 2333 CA Leiden, The Netherlands; phone: +31-71-5278467; fax: +33-71-527574; email: cacciato@strw.leidenuniv.nl
5. **Henk Hoekstra**: Leiden Observatory, Leiden University, Niels Bohrweg 2, 2333 CA Leiden, The Netherlands; phone: +31-71-5275594; fax: +33-71-5275743; email: hoekstra@strw.leidenuniv.nl
6. **Alina Kiessling**: NASA Jet Propulsion Laboratory, 4800 Oak Grove Drive, Pasadena, 91109 CA, USA; phone: +1-626-2981495; fax: +1-818-3548895; email: alina.a.kiessling@jpl.nasa.gov
7. **Donnacha Kirk**: Department of Physics & Astronomy, University College London, Gower Street, London WC1E 6BT, UK; phone: +44-79-977410621; fax: N/A; email: donnachakirk@gmail.com
8. **Thomas Kitching**: Mullard Space Science Laboratory, Holmbury St. Mary, Dorking, Surrey RH5 6NT, UK; phone: +44-1483-204218; fax: +44-1483-278312; email: t.kitching@ucl.ac.uk
9. **Rachel Mandelbaum**: Department of Physics, Carnegie Mellon University, 5000 Forbes Avenue, Pittsburgh, PA 15213, USA; phone: +1-412-2681714; fax: +1-412-6810648; email: rmandelb@andrew.cmu.edu
10. **Anais Rassat**: LASTRO, EPFL, Observatoire de Sauvigny, 1290 Versoix, Switzerland; phone: +33-671183365; fax: N/A; email: anais.rassat@epfl.ch
11. **Uros Seljak**: Department of Physics, Le Conte Hall, UC Berkeley, Berkeley CA 94708, USA; phone: +1-510-6662627; fax: N/A; email: useljak@berkeley.edu

Experts who potentially will contribute to the project:

- Ami Choi (Edinburgh)
- Alan Heavens (Imperial)
- Catherine Heymans (Edinburgh)
- Christopher Hirata (Ohio State)
- Adrienne Leonard (Saclay)
- Reiko Nakajima (Bonn)
- Peter Schneider (Bonn)

Benjamin Joachimi

Institute for Astronomy, University of Edinburgh

Main research interests:

- Nature of dark matter and dark energy; gravity on large scales
- Large-scale structure of Universe; galaxy formation and evolution
- Weak gravitational lensing; intrinsic alignment of galaxies
- Optimal design and exploitation of cosmological surveys
- Simultaneous analysis of cosmological probes and resulting synergies
- Spatial statistics and statistical inference

Education and career:

from Oct 13	Lecturer in Cosmology, University College London
Oct 12 – Sep 13	Ernest Rutherford fellow, University of Edinburgh
Oct 10 – Sep 12	DUEL/ Euclid fellow, University of Edinburgh
2010	PhD in Astrophysics, University of Bonn, <i>summa cum laude</i>
2007	Diploma in Physics, University of Bonn, <i>with distinction</i>

Awards, grants, distinctions:

2013	Royal Society Equipment Grant [1y; 15K £]
2012	STFC Ernest Rutherford Fellowship [5y; 530K £]
2012	Royal Astronomical Society Fellowship, <i>declined</i> [3y; 280K £]
2008	Prize 'Alumni-Preis der Universitäts-gesellschaft' (Bonn) [1K €]
2008	Stipendiary of the Wilhelm und Else Heraeus-Stiftung [1K €]
2007	BCGS 'honors programme' [3y; 10K €]
2007	Stipendiary of the Deutsche Telekom Stiftung [3y; 70K €]
2002	Fellow of the Studienstiftung des deutschen Volkes [5y; 5K €]

Relevant scientific responsibilities:

since 2012	Manager of Euclid Intrinsic Alignments tiger team
2010-2011	Leader of Euclid Intrinsic Alignments team (pre-selection)
since 2011	Work Package Manager for Euclid weak lensing data product validation
since 2011	Member of Euclid Galaxy Evolution Science Working Group
since 2008	Member of Euclid Weak Lensing Science Working Group
2013	Co-I of FP7 SPACE grant proposal on the PAU Survey
2012-2013	Main organizer (LOC & SOC) of DEX Workshops VIII & IX
since 2012	Member of KiDS (Kilo Degree Survey with VST)
since 2012	Member of the RCSLenS analysis team (Red Cluster Sequence Survey 2)

Selected relevant publications:

- **Joachimi, B.** & Schneider, P., '*The removal of shear-ellipticity correlations from the cosmic shear signal via nulling techniques*', *A&A* 488, 829 (2008)
- **Joachimi, B.** & Schneider, P., '*The removal of shear-ellipticity correlations from the cosmic shear signal: Influence of photometric redshift errors on the nulling technique*', *A&A* 507, 105 (2009)
- **Joachimi, B.** & Schneider, P., '*Intrinsic alignment boosting: Direct measurement of intrinsic alignments in cosmic shear data*', *A&A* 517, A4 (2010)
- **Joachimi, B.** & Bridle, S.L., '*Simultaneous measurement of cosmology and intrinsic alignments using joint cosmic shear and galaxy number density correlations*', *A&A* 523, A1 (2010)
- **Joachimi, B.**, Mandelbaum, R., Abdalla, F.B. & Bridle, S.L., '*Constraints on intrinsic alignment contamination of weak lensing surveys using the MegaZ-LRG sample*', *A&A*, 527, A26 (2011)
- **Joachimi, B.**, et al., '*Intrinsic galaxy shapes and alignments I: Measuring and modelling COSMOS intrinsic galaxy ellipticities*', *MNRAS*, *accepted* (2013)

Curriculum Vitae - Christopher Bonnett

Basic Information

Institut de Ciències de l'Espai, CSIC/IEEC
Fac de Ciències, Torre C5 par-2
Barcelona 08193
Spain

Telephone: +34 935 868042
e-mail : c.bonnett@gmail.com
Nationality: British/Belgian

Research Interests

cosmology, weak gravitational lensing, lensing systematic, galaxy bias, intrinsic alignments, surveys.

Academic Career

2011-present Marie Curie CosmoComp post-doc at CSIC/IEEC

23/09/2011 Phd, University Pierre et Marie Curie,FR supervisor Prof. Yannick Mellier

2007 MSc University of Leiden, NL

PARTICIPATION IN RESEARCH PROJECTS

Physics of the Accelerating Universe (PAU)

Lead of galaxy evolution sub-group

Euclid

Member of weak lensing science working group.

Member of Intrinsic Alignments working group.

Dark Energy Survey

Member of 2pt correlation and Covariance estimation group.

CFHTLenS

Lead of cross-correlation systematics group.

Curriculum Vitae – Michael L. Brown

Work address: Jodrell Bank Centre for Astrophysics
University of Manchester
Oxford Road
Manchester M13 9PL
United Kingdom

Nationality: Irish
Telephone: +44 161 3063913
E-mail: mbrown@jb.man.ac.uk

Research Interests: Observational cosmology, weak gravitational lensing, dark matter & dark energy, CMB polarization, early Universe physics.

Academic Career:

- 10/11 - present: Reader in Astrophysics, STFC Advanced Fellow
Jodrell Bank Centre for Astrophysics, University of Manchester.
- 10/06 - 09/11: Senior Research Associate.
Cavendish Laboratory and Kavli Institute for Cosmology,
University of Cambridge.
- 10/03 - 09/06: PPARC Postdoctoral Fellow.
Institute for Astronomy, University of Edinburgh.
- 03/03 - 09/03: Postdoctoral Research Assistant.
Institute for Astronomy, University of Edinburgh.
- 09/99 - 03/03: Ph.D. (Astronomy).
Institute for Astronomy, University of Edinburgh.
- 09/95 - 06/99: B.A. (Mod. Experimental Physics) (1st class, Gold Medal).
Trinity College, University of Dublin.

Major fellowships & awards gained as PI:

- European Research Council Starting Grant, 2012 - 2017.
- STFC Halliday Fellowship, 2011.
- STFC Advanced Fellowship, 2011 - 2016.
- PPARC Postdoctoral Fellowship, 2003 - 2006.

Service & professional activities:

- STFC Astronomy Advisory Committee, 2012 - present.
- SuperCLASS Weak Lensing leader & Steering Committee member.
- University of Manchester: PhD internal examiner.
- University of Cambridge: PhD internal and external examiner.
- Referee for MNRAS, ApJ & A&A journals, and STFC grant proposals.

Marcello Cacciato

Date of Birth: February 20, 1980
Place of Birth: Enna (Italy)
Nationality: Italian

Office 563, 2333 CA Oort Building
Leiden Observatory
The Netherlands
Email: cacciato@strw.leidenuniv.nl
URL: www.strw.leidenuniv.nl/~cacciato

EDUCATION

July 27, 2009 **Ph.D. in Astronomy (magna cum laude)**; Heidelberg University (Germany).
Thesis Title: Galaxy-Dark Matter Connection: from Astrophysics to Cosmology.
Advisor: Frank C. van den Bosch

RESEARCH EXPERIENCE

Sept 2012 - present: **Post-Doctoral Researcher**
Leiden Observatory, Leiden University, The Netherlands
October 2009-Aug 2012: **Post-Doctoral Researcher**
Racah Institute of Physics, The Hebrew University of Jerusalem, Israel

LIST OF SELECTED PUBLICATIONS

“A single model that describes weak lensing measurements around galaxies up to $z \sim 0.6$ ”,
Cacciato M., van Uitert E., Hoekstra H., soon to be submitted

“Cosmological Constraints from a Combination of Galaxy Clustering and Lensing -- III. Application to SDSS data”, **Cacciato M.**, van den Bosch F.C., More S., Mo H., Yang X., (arXiv:1207.0503)

“On combining galaxy clustering and weak lensing to unveil galaxy biasing via the halo model”,
Cacciato M., Lahav O., van den Bosch F., Hoekstra H., Dekel A., 2012, MNRAS, 426, 566

“Modeling Galaxy-Galaxy Weak Lensing with SDSS Groups” Li R., Mo H. J., Fan Z., **Cacciato M.**,
van den Bosch F. C., Yang X., More S., 2009, MNRAS, 394, 1016

“Galaxy Clustering & Galaxy-Galaxy Lensing: A Promising Union to Constrain Cosmological Parameters” **Cacciato M.**, van den Bosch F. C., More S., Li R., Mo H. J., Yang X., 2009, MNRAS, 394, 929

TEACHING EXPERIENCES

Semi-Analytical Models of Galaxy Formation (SAMs) [Summer Semester, 2010-11]
(Class series within ‘Advanced Cosmology’, Racah Institute of Physics, The Hebrew University)
The Halo Model of Large Scale Structure [Winter Semester, 2009-10]
(Class series within ‘Advanced Cosmology’, Racah Institute of Physics, The Hebrew University)

PROFESSIONAL SERVICE

Observatory Seminar organiser at Leiden Observatory [Since October 2012]
Referee for A&A, MNRAS, JCAP [Since March 2010]
Astrophysics seminar coordinator at Racah Institute of Physics [Since July 2010]

AWARDS

Rosenblum **Prize** 2010 (for the accomplished results, Hebrew University of Jerusalem, Israel)
Minerva **Fellowship** 2009-2011 (Max-Planck Society, Germany)

Curriculum Vitae - Henk Hoekstra

Basic Information:

Leiden Observatory Telephone: +31 (0)71 5275594
Leiden University Fax: +31 (0)71 5275743
Niels Bohrweg 2 e-mail: hoekstra@strw.leidenuniv.nl
2333 CA Leiden web: <http://www.strw.leidenuniv.nl/~hoekstra>
The Netherlands Nationality: Dutch

Research Interests

properties of clusters of galaxies- dark matter - dark energy - observational cosmology - gravitational lensing - surveys

Academic Career:

08/2008-present Assistant Professor, Leiden Observatory, Leiden University, NL
10/2004-07/2008 Assistant Professor, University of Victoria, Canada
09/2002-09/2004 Senior Research Associate, CITA, University of Toronto, Canada
10/2000-08/2002 Postdoctoral Fellow, CITA, University of Toronto, Canada
22/9/2000 PhD (cum laude), University of Groningen, NL
1995 MSc (cum laude), University of Groningen, NL

Honors, fellowship and scholarships:

ERC Starting Grant (2012-2017)
NWO Vidi grant (2009-2014)
Alfred P. Sloan Fellow (2007-2009)
Canadian Institute for Advanced Research Scholar (2004-present)
Beatrice D. Tremaine Fellowship (2001)

Service and Professional Activities

Euclid Cosmology Science Coordinator 2011-present
Euclid Weak Lensing Science Working Group Lead 2011-present
Science Advisory Committee ING telescopes 2009-present
Science Advisory Committee CFHT 2005-2008
Member Dutch Island Telescopes TAC 2009-2011
Member Canadian Time Allocation Committee 2005-2007

Curriculum Vitae – Alina Kiessling

Basic Information:

NASA Jet Propulsion Laboratory
4800 Oak Grove Drive
Pasadena, 91109
CA, USA

Telephone: +1 (626) 298-1495

email: Alina.A. Kiessling@jpl.nasa.gov

Nationality: German / Australian

Research Interests:

Gravitational Lensing: Weak – Cosmology: large scale structure of the Universe, dark matter, dark energy – Methods: N-body simulations, statistical

Academic Career:

01/2012 – present	Postdoctoral Fellow, NASA Jet Propulsion Laboratory, CA, USA
03/2011 – 09/2011	Postdoctoral Fellow, University of Edinburgh, Edinburgh, UK
31/03/2011	PhD, University of Edinburgh, Edinburgh, UK
2004	BSc (Space Science) Hons, LaTrobe University, Melbourne, Australia

Honors, fellowships and scholarships:

JPL Outstanding Postdoc Research Award for Astrophysics and Space Science, 2012
Marie Curie Early Stage Researcher Fellowship (2007 – 2010)
Menzies Center Bicentennial Scholarship (2006)
Australian Postgraduate Award (2005 – 2007)
Bok Prize runner up for the Most Outstanding Honors Thesis in Astronomy in Australia (2004)
LaTrobe University Prize for the Most Outstanding Student in Honors Physics (2004)

Service and Professional Activities:

NSF / AST Cosmology Panelist (2013)
Euclid Weak Lensing Cosmological Simulations Work Package Manager (2012 – present)
Euclid / LSST-DESC Covariance Matrices Task Coordinator / Member (2011 – present)
Euclid / LSST-DESC Intrinsic Alignments Tiger Team Member (2012 – present)
Astronomical Society of Australia Student Representative (2005 – 2007)

DONNACHA KIRK

Address Dept of Physics & Astronomy
University College London
Gower Street
WC1E 6BT
UK

Tel. 07977410621
Fax N/A
Email donnachakirk@gmail.com
Nationality Irish

Research interests

Gravitational lensing- Intrinsic Galaxy Alignment- Contamination forecasts- Combination of probes- Galaxy Surveys- Redshift Space Distortions- Modified Gravity- Photo-z Errors.

Academic Career

2012- PDRA, University College London
2011-12 PDRA, Imperial College London
2011 Post-doctoral Research Associate (PDRA), University College London
2008-11 PhD University College London
2006-07 MSci Experimental and Theoretical Physics, Cambridge, 1st
2003-06 BA Experimental and Theoretical Physics, Cambridge, 1st

Service & Professional Activities

Euclid Weak Lensing SG Member- Intrinsic Alignments sub-group	2011-present
EBEX Analysis Team Member	2012-present
MS-DESI Team Member- Joint Probes: WGL+LSS	2011-present
Dark Energy Survey, Theory & Combined Probes WG Member	2008-present
Dark Energy Survey, Weak Lensing WG Member- Intrinsic Alignments	2008-present
GREAT08 Shear Estimation Challenge, Core Group Member	2008-2010

Published Papers of Relevance

The cosmological impact of intrinsic alignment model choice for cosmic shear

D.Kirk, A.Rassat, O.Host & S.Bridle, MNRAS, 2012

Optimising cosmic shear surveys to measure modifications to gravity on cosmic scales

D.Kirk, I.Laszlo, S.Bridle & R.Bean, 2011

Disentangling dark energy and cosmic tests of gravity from weak lensing systematics

I.Laszlo, R.Bean, **D.Kirk** & S.Bridle, MNRAS, 2012

The impact of intrinsic alignments: cosmological constraints from a joint analysis of cosmic shear and galaxy survey data

D.Kirk, S.Bridle, & M.Schneider, MNRAS 2010

Thomas David Kitching

t.kitching@ucl.ac.uk, <http://www.thomaskitching.net>, +44 781 3135525, +44 1483 204 218

Career

November 2012 – Present	Lecturer Astrophysics	University College London, MSSL
October 2011 – Present	Royal Society University Research Fellow	Univeristy of Edinburgh, IfA
September 2011 – June 2012	Astronomical Statistics Lecturer	University of Edinburgh, IfA
January 2011 - October 2011	Royal Astronomical Society 2010 Fellow	University of Edinburgh, IfA
February 2009 - January 2011	Postdoctoral Euclid Research Fellow	University of Edinburgh, IfA
November 2006 - February 2009	Postdoctoral Research Assistant	University of Oxford, Physics
September 2007 - September 2008	Class A Lecturer in Astrophysics	St. Catherines College, Oxford

Summary of Awards

- Royal Astronomical Society Winton Capital Prize 2012
- Royal Society Univeristy Research Fellowship
- Royal Astronomical Society 2010 Fellowship (an inaugural award of this Fellowship)
- Outstanding Contribution, salary increment award, School of Physics Univeristy of Edinburgh 2010

Education

PhD, Cosmology, PPARC funded; Thesis : Constraining the equation of state of dark energy using 3D weak lensing.
MSci 1st Class (Hons), Physics, *Imperial College London*

Grants (relevant for Euclid)

- Jan 2011, Co-I STFC Euclid funding for implementation phase activities (PI: M. Cropper)
- Feb 2010, Co-I STFC Euclid funding for definition phase activities (PI: M. Cropper)
- Jan 2010, €10,985 – PI PASCAL Network (funded by the EC) for GREAT10 PASCAL Challenge
- May 2008, £50 K - Co-I STFC further bridging support for Euclid satellite project (PI: M. Cropper)
- Mar 2008, €21,500 – Co-I PASCAL Network (funded by the EC) for GREAT08 PASCAL Challenge (PI: S. Bridle)

PhD Supervisor Roles

- 2011 – 2012 PhD Co-supervisor Marika Asgari (3D gravitational lensing)
- 2010 – 2012 PhD Co-supervisor David Harvey (dark matter cluster lensing constraints)
- 2009 – 2011 PhD Co-supervisor Emma Grocutt (weak lensing tomography)

Key Roles

<i>Euclid – ESA Cosmic Visions</i>	<ul style="list-style-type: none">• Euclid Consortium Cosmology Science Coordinator - 2011 to Present• Euclid Weak Lensing Coordinator – 2011 to Present
<i>GREAT Simulations (GRavitational lEnsing Accuracy Testing)</i>	<ul style="list-style-type: none">• PI of the Mapping Dark Matter challenge• PI of the GREAT10 simulations challenge
<i>Pan-STARRS</i>	<ul style="list-style-type: none">• Lead Data Reduction and Verification Group (DRAVG), Edinburgh – 2009 to 2011• Core member of Key Project 11 (cosmological lensing)
<i>CFHTLenS (CFHTLS Lensing Survey)</i>	<ul style="list-style-type: none">• Core team member• Coordinator Cosmology Working Group

Publication Summary

- 68 articles since 2005; (ADS) H-index = 20. Total Citations = 1231. Top Citation = 127.

Rachel Mandelbaum

Carnegie Mellon University, Dept. of Physics
5000 Forbes Ave.
Pittsburgh, PA 15213, USA
412-268-1714 ♦ rmandelb@andrew.cmu.edu

Research interests:

Observational cosmology, dark matter, dark energy, gravitational lensing, galaxy clustering, galaxy formation and evolution, galaxy cluster formation and evolution, large-scale structure

Education

Ph. D., Physics, Princeton University, 8/2006

A. B., Physics, Princeton University, 6/2000

Professional Experience

9/2011-present: Assistant Professor, Carnegie Mellon University, Dept. of Physics

11/2009-6/2012: Associate Research Scholar, Princeton Univ., Dept. of Astrophysical Sciences

9/2006-11/2009: Hubble Fellow, Institute for Advanced Study, Astrophysics

Honors and awards

Alfred P. Sloan Fellow (2013-)

US Department of Energy Early Career Award (2012-2017)

Annie Jump Cannon Prize of the American Astronomical Society (2011)

Hubble Fellow (2006-2009)

Relevant Professional Activities

Co-leader, GREAT3, the upcoming weak lensing community data challenge (late 2011-present)

Co-leader, LSST Dark Energy Science Collaboration weak lensing working group (June 2012-present)

Co-leader, weak lensing working group for the Hyper-SuprimeCam (HSC) survey

Co-organizer of workshop (summer 2013) on weak lensing at the Aspen Center for Physics

Selected Publications Relevant to this Proposal

A. Smargon, R. Mandelbaum, N. Bahcall, M. Niederste-Ostholt, *Detection of intrinsic cluster alignments to 100 Mpc/h in the Sloan Digital Sky Survey*, 2012, MNRAS, 423, 856.

J. Blazek, R. Mandelbaum, U. Seljak, R. Nakajima, *Separating intrinsic alignment and galaxy-galaxy lensing*, 2012, JCAP, 5, 41.

B. Joachimi, R. Mandelbaum, F. B. Abdalla, S. L. Bridle, *Constraints on intrinsic alignment contamination of weak lensing surveys using the MegaZ-LRG sample*, 2011, A&A, 527, 26.

R. Mandelbaum, C. Blake, S. Bridle, F. B. Abdalla, et al., *The WiggleZ Dark Energy Survey: Direct constraints on blue galaxy intrinsic alignments at intermediate redshifts*, 2011, MNRAS, 410, 844.

C. M. Hirata, R. Mandelbaum, et al., *Intrinsic galaxy from the 2SLAQ and SDSS surveys: luminosity and redshift scalings and implications for weak lensing surveys*, 2007, MNRAS, 381, 1197.

R. Mandelbaum, C. Hirata, M. Ishak, U. Seljak, J. Brinkmann, *Detection of large-scale intrinsic ellipticity-density correlation from the Sloan Digital Sky Survey and implications for weak lensing surveys*, 2006, MNRAS, 367, 611.

Anais Rassat, Ph.D.

LASTRO, EPFL, Observatoire de Sauverny, 1290 Versoix, Switzerland

Tel (French mobile/cell): +33 (0)6 71 18 33 65

anais.rassat@epfl.ch

<http://anaisrassat.tumblr.com>

Nationality: French

Summary & Research Interests

Euclid member and long-term postdoctoral researcher in observational cosmology

Research Interests: large scale structure, weak & strong lensing, intrinsic alignments, probe combination

Professional Experience

Sept 2011- present: Laboratory of Astrophysics (LASTRO), EPFL. Long-Term (5 year) Postdoctoral Scientist

Nov 2007 - Aug 2011: Service d'Astrophysique, AIM, CEA-CNRS. Postdoctoral Scientist / Engineer (4 years)

Education

2007: Ph.D. Observational Cosmology, University College London, UK

2003: Masters of Advanced Studies in Mathematics (Part III Maths), University of Cambridge, UK

2002: BSc Maths & Physics with Astrophysics, King's College, London, UK

Academic Prizes

2004-2007: Perren Fund Studentship (Ph.D.)

2002: Institute of Mathematics and its Applications Prize, King's College, London

2001: Florence Hughes Prize, King's College, London

Professional Services

- since 2011: Deputy lead of Euclid Consortium COMS (communications) group.
- since 2011: Lead Education and Public Outreach (EPO) scientist for the Euclid Consortium.
- 2010-2011: Lead of Euclid Consortium COMS group.
- 2007-2010: Co-coordinator of Euclid Imaging Consortium BAO working group.
- Presented 23 talks at numerous international conferences and workshops, 7 times invited speaker, 4 times session chair and 14 times seminar speaker.
- Twice External Examiner for Ph.D. defense of P. Arnalte-Mur (2011, Prof. V. Martinez, Valencia, Spain), and A. Labatie (Dec. 2012, Prof. J.-L. Starck, AIM, CEA, France)
- Supervision of 3 Masters students: R. Harrison (2012, → Imperial, Masters), F. Lanusse (2011, → Ph.D. Saclay with J.-L. Starck), B. Leistedt (2010, → Ph.D. UCL with H. Peiris)
- Referee for MNRAS (Monthly Notices of Royal Astronomical Society) and IJMP (International Journal of Modern Physics)

Recent Invited Talks

- **Invited Lecturer at Euclid-ESA School** for ESA-Engineers, IAP, France, 'Combining Probes, Fisher Matrices and Statistics', Sept. 2012
- **'Spherical 3D Analysis of Cosmological Galaxy Surveys'**, UNIGE, Switzerland, March 2012
- **'Reconstructing the Integrated Sachs-Wolfe effect'**, ADA VI Conference, Tunisia, May 2010

Publications

14 refereed publications, including 3 first author, 5 second author papers.

Total Citations: 644 (ADS March 2013)

Co-author of 4 public codes: [iCosmo](#), [3DEX](#), [MRS3D](#), [StatAnomalies](#)

Editor and Co-Author of Euclid Imaging Consortium Science Book ('Yellow Book', 2010)

Co-Author of Euclid Definition Study Report ('Red Book', 2012)

Curriculum Vitae

Uroš Seljak

University of California at Berkeley
Department of Physics
441 Le Conte Hall
Berkeley CA 94720
email: useljak@berkeley.edu

Research Interests

Cosmology and Theoretical Astrophysics

Research Experience

7/2006-present: professor, University of California at Berkeley, Berkeley, USA.

7/2006-8/2013: professor, ITP, Zurich University, Zurich Switzerland.

2004-2006: research staff, ICTP, Trieste, Italy.

1999-2007: assistant professor, Department of Physics, Princeton University, Princeton, USA.

1995-1998: Smithsonian Fellow, Harvard-Smithsonian Center For Astrophysics, Cambridge, USA.

Education

June 1995 : PhD in Physics, MIT (advisor: E. Bertschinger)

July 1991 : MS in Physics, Univ. of Ljubljana, Slovenia.

October 1989 : BS in Physics, Univ. of Ljubljana, Slovenia.

Awards

2002: NSF CAREER grant award

2001: Alfred P. Sloan Fellow

2001: Helen B. Warner award of American Astronomical Society

2000: David and Lucille Packard Fellow

2000: Princeton 250th Anniversary Fund Award for Innovation in Undergraduate Education

1993 : Alan H. Barrett Prize for Excellence in Astrophysics at MIT.

Supervisory Experience: 14 graduate students, 10 postdocs

Publications

180 refereed papers, 197 total papers, 23000 citations (H=68), 5600 normalized citations (H=37) (NASA ADS)