



UiO :



Roseland  
Centre  
for Solar  
Physics

# Long-period intensity pulsations in coronal loops

**Clara Froment**

Roseland Centre for Solar Physics - University of Oslo


P. Antolin, V. Henriques & L. Rouppe van der Voort  
& ISSI team on coronal loops

<http://www.issibern.ch/teams/observecoronloop/>

BUKS 2018 - 3-7 September 2018

# Long-period EUV pulsations are very common




There are pulsations with several hours of periods in there!





# Long-period EUV pulsations are very common

There are pulsations with several hours of periods in there!



# Pulsating loops in almost every active region

➔ Intensity pulsations (2 - 16 hrs) in coronal EUV channels

On-disk detections:

- 917 events found in 13 yrs of EIT (195 Å)  
*Auchère et al. 2014*




54% AR, 45% QS

- 3181 events found in 6 yrs of AIA  
*Froment 2016, PhD thesis*

67% AR, 33% QS

Off-limb detections:

- 2981 events found in 7 yrs of AIA off-limb  
62% AR, 38% QS



Frequency: 31.3  $\mu\text{Hz}$   
Frequency width: 9.2  $\mu\text{Hz}$   
Frequency resolution: 4.6  $\mu\text{Hz}$   
Period: 8.9 h  
Maximum power: 37.6  $\sigma$   
Average power: 21.7  $\sigma$   
Average intensity: 118.4  
Relative amplitude: 0.196  
Threshold: 10.0  $\sigma$   
Theta: 109.8°  
Rho: 1070.5 arcsec  
Surface: 188.0 pixels



# Pulsating loops in almost every active region

➔ **Intensity pulsations** (2 - 16 hrs) in coronal EUV channels

On-disk detections:

- **917 events found in 13 yrs of EIT (195 Å)**  
*Auchère et al. 2014*

54% AR, 45% QS




- **3181 events found in 6 yrs of AIA**  
*Froment 2016, PhD thesis*

67% AR, 33% QS

Off-limb detections:

- **2981 events found in 7 yrs of AIA off-limb**  
62% AR, 38% QS

**Database and full statistics soon publicly available**  
*Froment et al., in prep*



**Significance critically re-assessed in *Auchère et al. 2016a***

Wavelet analysis with the Torrence & Compo 1998 code


➔ how to use the TC98 code with **any noise model** instead of the built-in white or red noise, and with **global confidence levels**

Surface: 188.0 pixels

# What are these pulsations?

## ✗ Not triggered by an other event, Not a mechanical oscillation

- ➔ Not connected to late phase of flare or CMEs
- ➔ Not the signature of vibration mode but a periodic train of pulses of random amplitudes (*Auchère et al. 2016*)




## ✓ Thermal phenomenon

- ➔ Current explanation for these pulsations:  
**Coronal counterpart of thermal nonequilibrium cycles**
- ➔ Implications for coronal heating: **spatial location and timescale**

# Evaporation/condensation cycles in loops


## Detection with AIA

9 hr pulsations in AR 11499




## AIA observations

Froment et al. 2015



## Synthetic intensities from a 1D hydrodynamic simulation



Froment et al. 2017 & 2018



- ➔ **Periodic rain event observed for the first time (with SDO/AIA, Auchère et al. 2018)**
- ➔ **How cold can it get?**
- ➔ **At least some events should show coronal rain down to chromospheric temperatures**



# Multi-thermal analysis off-limb with SDO and SST



*Froment, Antolin, Henriques & Rouppe van der Voort, in prep*

- ➔ Period of ~6.h in almost all the channels
- ➔ Swedish 1-m Solar Telescope (SST) observations at **one footpoint** during the **cooling phase of one of the cycles**


# SST observations for one cooling phase

- ➔ Observation of the cycle from coronal to **chromospheric temperature**
- ➔ High-resolution coronal rain observations

SDO/AIA 171

SDO/AIA 304

SST/CHROMIS Ca II K + 0.59 Å




## SST data:

~30 min during of the cooling phase  
At the middle of the AIA sequence

- **CRISP: H $\alpha$  (6563 Å)**  
pixel size: 0.06''
- **CHROMIS: Ca II K (3934 Å)**  
pixel size: 0.04''

# SST observations for one cooling phase

- ➔ Observation of the cycle from coronal to **chromospheric temperature**
- ➔ High-resolution coronal rain observations






# Evolution of the temperature and the density

## Analysis of the thermal structure

➔ **Reconstruction of the Differential Emission Measure (DEM)** - code from *Cheung et al, 2015*




- ➔ **Cycles** ( $\sim 6\text{h}$ ) in the DEM-weighted temperature and the total EM ( $\propto n_e^2$ )
- ➔ **The temperature increases always before the total EM**
- ➔ Temperature and width anti-correlated  
**Thermal width increase**  
➔ **cooling phases**

- ➔ **Same conclusions as for on-disk observations of pulsating loops**
- ➔ **Strong evidence of TNE**

# Observation of cooling with SDO/AIA

**Time lags** (same technique as in *Viall & Klimchuk, 2012*):

- peak **cross-correlation** values (pairs of channels)
- to **reconstruct the order of the channels** and thus **the temperatures**




- ➔ **Widespread cooling, same patterns of time lags as on-disk observations**
- ➔ **The pulsating loops have the same cooling behavior as the rest of the active region**

# Thermodynamic of the rain

SDO/AIA 171 Å  
 SST/CRISP H-alpha + 0.70 Å  
 SST/CHROMIS Ca II K + 0.59 Å

- Intensity threshold to detect rain pixels
- Stack of N slabs in time




- Gaussian fit of the H $\alpha$  condensation profiles

$$FWHM = 2\sqrt{2 \ln 2} \frac{\lambda_0}{c} \sqrt{\frac{2k_B T}{m_H} + \cancel{\nu_{mic}^2}}$$

➔ upper bounds for the plasma temperature



# Thermodynamic of the rain



## ➔ Preliminary results:

*Average temperature: ~25 000 K*

*Average Doppler velocity: ~39 km/s*

*Projected velocities ~70 km/s*

*Total velocity ~80 km/s*

## ➔ Temperature and velocities consistent with other rain studies

*(Antolin & Rouppe Van der Voort 2011, Ahn et al. 2014, Antolin et al. 2015)*

# Conclusions

- **Long-period intensity pulsations** (several hours) are **very common in coronal loops**
- **Long-period intensity pulsations** are the coronal counterpart of **thermal nonequilibrium cycles** and thus of **quasi-constant** and **highly stratified heating**
- The pulsating loops studied with AIA off-limb show the same thermal behaviour as for on-disk pulsating loops
- These observations allow us to **probe the bulk of the cooling phases** and emphasise that these **pulsations and coronal rain are two aspects of the same phenomenon**
- **Implication for circulation of mass and energy in the solar corona**

# To go further

Some open questions:

- ▶ What fraction of the coronal volume experiences TNE?
- ▶ Are the non-pulsating loops and diffuse emission produced by a completely different heating deposition in time and space?
- ▶ What determines whether a condensation forms or the thermal collapse is aborted before reaching chromospheric temperatures?

ISSI team selected in 2017:  
Observed Multi-Scale Variability of Coronal Loops as  
a Probe of Coronal Heating  
<http://www.issibern.ch/teams/observecoronloop/>